

THE 67<sup>TH</sup> INTERNATIONAL



# OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

BOOK OF  
ABSTRACTS

2024



Vilnius  
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys  
Rimantas Naina  
Vincentas Adomaitis  
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press  
9 Saulėtekio Av., III Building, LT-10222 Vilnius  
info@leidykla.vu.lt, [www.leidykla.vu.lt/en/](http://www.leidykla.vu.lt/en/)  
[www.knygynas.vu.lt](http://www.knygynas.vu.lt), [www.journals.vu.lt](http://www.journals.vu.lt)

Bibliographic information is available  
on the Lithuanian Integral Library Information System (LIBIS) portal [www.ibiblioteka.lt](http://www.ibiblioteka.lt)  
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

<b>FORMATION AND INVESTIGATION OF BIPOLAR AQUEOUS SODIUM ION BATTERIES</b> . . . . .	<b>291</b>
Dovilė Škarnulytė, Milda Petrulevičienė, Jurga Juodkazytė, Linas Vilčiauskas	
<b>SYNTHESIS AND LUMINESCENT PROPERTIES OF EU-DOPED STRONTIUM CHLORAPATITE</b> . . . . .	<b>292</b>
Simona Bendziute, Inga Grigoraviciute, Arturas Katelnikovas, Aleksej Zarkov	
<b>STRUCTURAL AND NMR PROPERTIES OF SUPRAMOLECULAR COMPLEXES OF DRUG MOLECULES</b> . . . . .	<b>293</b>
Benjaminas Malmiga, Kęstutis Aidas	
<b>SYNTHESIS OF TARGETED CONDENSED THIOIMIDAZOLES</b> . . . . .	<b>294</b>
Martyna Paulauskaitė, Gintarė Sapežinskaitė, Indrė Misiūnaitė, Ieva Žutautė	
<b>P3: Biochemistry, Biophysics and Biotechnology; Biology, Genetics and Biomedical Sciences</b>	
<b>CAPDROP: A NOVEL METHOD FOR PERIPHERAL BLOOD scRNA-seq</b> . . . . .	<b>295</b>
Emilė Pranauskaitė, Linas Mažutis	
<b>COMPARATIVE STUDIES OF THE ANTIOXIDANT PROPERTIES OF DIOSMIN AND QUERCETIN IN THE MODEL SYSTEM OF DOPAMINE OXIDATION</b> . . . . .	<b>296</b>
Iryna Povshedna, Vladyslav Udovytskyi, Iryna Pashchenko, Viktoriia Lyzhniuk, Vadym Lisovyi, Volodymyr Bessarabov, Andriy Goy	
<b>INHIBITION OF PERIODONTAL DISEASES SPECIFIC MIRNAS: NEW THERAPEUTIC APPROACH</b> . . . . .	<b>297</b>
Gabrielė Židonytė, Benita Buragaitė-Staponkienė, Kristina Šnipaitienė, Adomas Rovas, Alina Purienė, Raminta Vaičiulevičiūtė, Eiva Bernotienė, Sonata Jarmalaitė	
<b>INVESTIGATION OF ANXA4 FUNCTION IN CELL PLASMA MEMBRANE PERMEABILIZATION, RESEALING AND CELL VIABILITY POST-ELECTROPORATION</b> . . . . .	<b>298</b>
Dominykas Makarovas, Baltramiejus Jakštys, Saulius Šatkauskas	
<b>CREATION OF MUTANT VARIANT K102R OF YEAST SACCHAROMYCES CEREVISIAE GENE SUP35</b> . . . . .	<b>299</b>
Ieva Jankauskaitė, Justina Versockienė, Eglė Lastauskienė	
<b>MINIMIZING CHEMOTHERAPY SIDE EFFECTS: CALCIUM SONOPORATION</b> . . . . .	<b>300</b>
Reda Rulinskaitė, Rūta Palepšienė, Saulius Šatkauskas, Renaldas Raišutis, Martynas Maciulevičius	
<b>EXPLORING THE IS200/IS605 FAMILY TRANSPOSABLE ELEMENTS FOR NOVEL GENOME EDITING TOOLS</b> . . . . .	<b>301</b>
Gytis Druteika, Tautvydas Karvelis	
<b>STRUCTURAL VARIABILITY OF PRION PROTEIN AMYLOID FIBRILS DEPENDS ON AGITATION INTENSITY</b> . . . . .	<b>302</b>
Kamile Mikalauskaite, Mantas Ziaunys, Vytautas Smirnovas	
<b>CRISPR-CAS9 GENOME ENGINEERING IN <i>KLUYVEROMYCES MARXIANUS</i> FOR ENHANCED SECRETION OF RECOMBINANT ANTIBODIES</b> . . . . .	<b>303</b>
Justina Žičkutė, Danguolė Žiogienė, Alma Gedvilaitė	

## COMPARATIVE STUDIES OF THE ANTIOXIDANT PROPERTIES OF DIOSMIN AND QUERCETIN IN THE MODEL SYSTEM OF DOPAMINE OXIDATION

Iryna Povshedna<sup>1</sup>, Vladyslav Udovytskyi<sup>1</sup>, Iryna Pashchenko<sup>1</sup>, Viktoriia Lyzhniuk<sup>1</sup>, Vadym Lisovyi<sup>1,2</sup>, Volodymyr Bessarabov<sup>1</sup>, Andriy Goy<sup>1</sup>

<sup>1</sup>Department of Industrial Pharmacy, Kyiv National University of Technologies and Design, Ukraine

<sup>2</sup>Department of Chemical Technology and Resource Saving, Kyiv National University of Technologies and Design, Ukraine  
[povshedna.io@knuud.edu.ua](mailto:povshedna.io@knuud.edu.ua)

Parkinson's disease (PD) is a common neurodegenerative disorder. It is associated with the degeneration of dopaminergic neurons and is characterized by tremor, postural instability, rigidity, and hypokinesia. The treatment of PD primarily involves symptomatic therapy, which includes administering levodopa, which is converted to dopamine in the brain [1]. However, treating PD with levodopa is problematic because both levodopa and its metabolite dopamine can be oxidised, usually by oxidative stress. Therefore, additional administration of antioxidants is needed to prevent oxidation of levodopa and dopamine.

Flavonoids are a well-known group of antioxidants, which are biologically active phenolic substances found in various plants. Recent studies have shown that some flavonoids can protect dopaminergic neurons by inhibiting dopamine oxidation and modulating antioxidant signaling pathways [2-3].

The study examined the effect of two well-known bioflavonoids, diosmin and quercetin, on the rate of dopamine oxidation and compared their antioxidant activity in this system. Kinetic studies were conducted using the spectrophotometric method, which recorded the increase in the optical absorption of the reaction mixture over time at a wavelength of 500 nm. The rate of dopamine oxidation was determined by calculating the first-order reaction rate constant.

Analyzing the dependence of the first-order reaction rate constant of dopamine oxidation on the concentration of diosmin in the system, it can be argued that a dose-dependent effect of oxidation inhibition is observed. When diosmin was added to the system at a concentration of 25  $\mu\text{M}$ , the rate of dopamine oxidation significantly decreased by 3.07 times:  $K_{H(0)}^1 = (3.90 \pm 0.03) \times 10^{-3}$  1/s and  $K_{H(25)}^1 = (1.27 \pm 0.04) \times 10^{-3}$  1/s ( $p \leq 0,05$ ); with an increase in concentration to 50  $\mu\text{M}$  the rate of dopamine oxidation decreased by 4.02 times:  $K_{H(50)}^1 = (0.97 \pm 0.08) \times 10^{-3}$  1/s, and at a concentration of 100  $\mu\text{M}$  – 5.57 times:  $K_{H(100)}^1 = (0.70 \pm 0.05) \times 10^{-3}$  1/s.

Instead, when quercetin was added to the system at a concentration of 25  $\mu\text{M}$  the rate of dopamine oxidation decreased significantly by only 1.35 times:  $K_{H(0)}^1 = (3.90 \pm 0.03) \times 10^{-3}$  1/s and  $K_{H(25)}^1 = (2.88 \pm 0.04) \times 10^{-3}$  1/s ( $p \leq 0,05$ ). When the quercetin concentration is increased to 50 and 100  $\mu\text{M}$  the oxidation rate decreases by 1.65 and 3.02 times, respectively:  $K_{H(50)}^1 = (2.36 \pm 0.06) \times 10^{-3}$  1/s and  $K_{H(100)}^1 = (1.29 \pm 0.03) \times 10^{-3}$  1/s.

It has been established that the bioflavonoids diosmin and quercetin exhibit antioxidant properties in relation to dopamine oxidation. However, diosmin is a more effective inhibitor of dopamine oxidation. Therefore, it is advisable to prefer diosmin when modelling a pharmaceutical composition for use in combination therapy with levodopa drugs for the treatment of Parkinson's disease.

[1] E. K. J. Pauwels, G. J. Boer. Parkinson's Disease: A Tale of Many Players. Medical principles and practice : international journal of the Kuwait University, Health Science Centre, 32(3), 155–165 (2023). <https://doi.org/10.1159/000531422>

[2] S. L. Hor, S. L. Teoh, W. L. Lim. Plant Polyphenols as Neuroprotective Agents in Parkinson's Disease Targeting Oxidative Stress. Current drug targets, 21(5), 458–476 (2020). <https://doi.org/10.2174/1389450120666191017120505>

[3] Z. D. Zhou; S. P. Xie, W. T. Saw, P.G.H. Ho, H.Y. Wang, L. Zhou, Y. Zhao, E.K. Tan. The Therapeutic Implications of Tea Polyphenols against Dopamine (DA) Neuron Degeneration in Parkinson's Disease (PD). Cells, 8, 911 (2019). <https://doi.org/10.3390/cells8080911>